

Extended Hamiltonian Formalism of the Pure Space-Like Axial Gauge Schwinger Model

Yuji NAKAWAKI and Gary McCARTOR*

Division of Physics and Mathematics, Faculty of Engineering

Setsunan University, Osaka 572-8508, Japan

**Department of Physics, Southern Methodist University*

Dallas TX 75275, USA

Abstract

We demonstrate that pure space-like axial gauge quantizations of gauge fields can be constructed in ways which are free from infrared divergences. To do so we must extend the Hamiltonian formalism to include residual gauge fields. We construct an operator solution and an extended Hamiltonian of the pure space-like axial gauge Schwinger model. We begin by constructing an axial gauge formulation in auxiliary coordinates: $x^\mu = (x^+, x^-)$, where $x^+ = x^0 \sin \theta + x^1 \cos \theta$, $x^- = x^0 \cos \theta - x^1 \sin \theta$ and we take $A_- = A^0 \cos \theta + A^1 \sin \theta = 0$ as the gauge fixing condition. In the region $0 \leq \theta < \frac{\pi}{4}$, we can take x^- as the evolution parameter and construct a traditional canonical formulation of the temporal gauge Schwinger model in which residual gauge fields dependent only on x^+ are static canonical variables. Then we extrapolate the temporal gauge operator solution into the axial region, $\frac{\pi}{4} < \theta < \frac{\pi}{2}$, where x^+ is taken as the evolution parameter. In the axial region we find that we have to change representations of the residual gauge fields from one realizing the PV prescription to one realizing the ML prescription in order for the infrared divergences resulting from $(\partial_-)^{-1}$ to be canceled by corresponding ones resulting from the inverse of the hyperbolic Laplace operator. We overcome the difficulty of constructing the Hamiltonian for the residual gauge fields by employing McCartor and Robertoson's method, which gives us a term integrated over $x^- = \text{constant}$. Finally, by taking the limit $\theta \rightarrow \frac{\pi}{2} - 0$ we obtain an operator solution and the Hamiltonian of the axial gauge (Coulomb gauge) Schwinger model in ordinary coordinates. That solution includes auxiliary fields and the representation space is of indefinite metric, providing further evidence that “physical” gauges are no more physical than “unphysical” gauges.

§1. Introduction

Axial gauges, $n^\mu A_\mu = 0$, specified by a constant vector n^μ , have been used recently in spite of their lack of manifest Lorentz covariance. One reason is that the Faddeev-Popov ghosts decouple from the theory in the axial gauge formulations.¹⁾ The case of $n^2 = 0$, the light-cone gauge, has been extensively used in light-front field theory (LFFT) in attempts to find nonperturbative solutions of QCD. In LFFT one usually makes the change of variables, $x_l^+ = \frac{x^0+x^3}{\sqrt{2}}$, $x_l^- = \frac{x^0-x^3}{\sqrt{2}}$,²⁾ and specifies quantization conditions on the hyperplane $x_l^+ = 0$. Advantages are that the vacuum state contains only of particles with nonnegative longitudinal momentum and that there exist relativistic bound-state equations of the Schrödinger-type. (For a good overview of LFFT see Ref. 3).)

It has been known for some time that axial gauge formulations are not ghost free, contrary to what was originally expected and is still sometimes claimed. It was first pointed out by Nakanishi⁴⁾ that there exists an intrinsic difficulty in the axial gauge formulations so that an indefinite metric is indispensable even in QED. It was also noticed that in order to bring perturbative calculations done in the light-cone gauge into agreement with calculations done in covariant gauges, spurious singularities of the free gauge field propagator have to be regularized not as principal values (PV), but according to the Mandelstam-Leibbrandt (ML) prescription⁵⁾. Shortly afterwards, Bassetto et al.⁶⁾ obtained the ML form of the propagator in a canonical formalism. They quantized at equal time in light-cone gauge, which required the introduction of a Lagrange multiplier field and its conjugate. These may be viewed as residual gauge degrees of freedom. Furthermore, Morara and Soldati⁷⁾ found just recently that the same is true in the light-cone temporal gauge formulation in which x_l^+ is again taken as the evolution parameter but $\frac{A_0+A_3}{\sqrt{2}} = 0$ is taken as the gauge fixing condition. McCartor and Robertson⁸⁾ showed that in the light-cone axial gauge formulation, where $\frac{A_0-A_3}{\sqrt{2}} = 0$ is taken as the gauge fixing condition, the translational generator P_+ consists of physical degrees of freedom integrated over the hyperplane $x_l^+ = \text{constant}$ and residual gauge degrees of freedom integrated over the hyperplane $x_l^- = \text{constant}$.

In McCartor and Robertson's work the residual degrees of freedom were viewed as integration constants necessary to completely specify the solution to the constraint equations which relate constrained degrees of freedom to the independent degrees of freedom. That is the attitude we will adopt in the present paper. Such equations often occur in quantum field theory and little attention has been given to the question of what boundary conditions are appropriate to completely specify their solution. In the present paper we will attend to that question. An example of the type of equation we have in mind is the equation for A_+

in the light-cone gauge

$$\partial_-^2 A_- + \partial_- \partial_i A^i = J_- . \quad (1.1)$$

This equation admits a solution of the form

$$C(x^+, \mathbf{x}^\perp) + x^- B(x^+, \mathbf{x}^\perp) + F(x),$$

where $F(x)$ is any solution to the equation. McCartor and Robertson showed that if $F(x)$ is taken to be the usual solution, which results from the replacement $\partial_\mu \rightarrow ik_\mu$ in the mode expansion, then neither $C(x^+, \mathbf{x}^\perp)$ or $B(x^+, \mathbf{x}^\perp)$ are zero in the case of QED. On the other hand, in an earlier work⁹⁾, we showed that for the Schwinger model in the light-cone gauge, $B(x^+, \mathbf{x}^\perp)$ is zero while $C(x^+, \mathbf{x}^\perp)$ is not. It will be seen below that it is this last case that is peculiar: for all the solutions we shall give, both $B(x^+, \mathbf{x}^\perp)$ and $C(x^+, \mathbf{x}^\perp)$ are nonzero except for the special case of the light-cone gauge where $B(x^+, \mathbf{x}^\perp)$ happens to vanish.

Because the axial gauges can be viewed as continuous deformations of the light-cone gauge, extensions of the ML prescription outside the light-cone gauge formulation have also been studied. Lazzizzera¹⁰⁾, and Landshoff and Nieuwenhuizen¹¹⁾, constructed canonical formulations for the non-pure space-like case ($n^0 \neq 0, n^2 < 0$) in ordinary coordinates. However, in spite of many attempts, no one has succeeded in constructing a consistent, pure space-like axial gauge ($n^0 = 0, n^2 < 0$) formulation. One problem is that when the residual gauge fields are introduced as integration constants that occur when one integrates the axial gauge constraints, one cannot obtain their quantization conditions from the Dirac quantization procedure¹²⁾. Another problem is that the residual gauge fields are independent of x^3 in the gauge $A_3 = 0$ so that the Hamiltonian cannot be obtained by integrating densities made of those residual gauge fields over the three dimensional hyperplane $x^0 = \text{constant}$.

This motivates us to study the problem of introducing the residual gauge fields into pure space-like axial gauge formulations. To construct the Hamiltonians and to determine quantization conditions, we construct the pure space-like gauge as a continuous deformation of the light-cone gauge. In a previous, preliminary work¹³⁾ we have constructed an axial gauge formulation of noninteracting abelian gauge fields in the auxiliary coordinates $x^\mu = (x^+, x^-, x^1, x^2)$, where

$$x^+ = x^0 \sin\theta + x^3 \cos\theta, \quad x^- = x^0 \cos\theta - x^3 \sin\theta \quad (1.2)$$

and $A_- = A^0 \cos\theta + A^3 \sin\theta = 0$ is taken as the gauge fixing condition. The same framework was used previously by others to analyze two-dimensional models.¹⁴⁾ In the region $0 \leq \theta < \frac{\pi}{4}$ we take x^- as the evolution parameter and construct the canonical temporal gauge formulation; in that case the residual gauge fields are static canonical variables. By continuation,

we can obtain operators satisfying the field equations in the axial region $\frac{\pi}{4} < \theta < \frac{\pi}{2}$, where x^+ is taken as the evolution parameter. In the axial region we cannot use the traditional way of constructing the Hamiltonian for the residual gauge fields; we obtain it by integrating the divergence equation of the energy-momentum tensor over a suitable closed surface. Because the residual gauge fields do not depend on x^- , their boundary surface ($x^- \rightarrow \pm\infty$) contributions have to be kept. As a consequence, we obtain the Hamiltonian which includes a part from integrating a density involving the residual gauge fields over $x^- = \text{constant}$. Then, by taking the limit $\theta \rightarrow \frac{\pi}{2} - 0$ we obtain the Hamiltonian in the axial gauge $A_3 = 0$ formulation in ordinary coordinates.

In this paper we proceed to interacting gauge fields. As a first step in finding ways to introduce the residual gauge fields into interacting axial gauge theories, we consider the exactly solvable Schwinger model. Previous attempts to construct the axial gauge ($A_1 = 0$) operator solution to the Schwinger model have used a representation space isomorphic to (copies of) that of a positive metric free massive scalar field $\tilde{\Sigma}$. In other words, they have assumed that ($A_1 = 0$) is a physical gauge in that the entire representation space is physical. All such attempts encounter severe infrared difficulties.¹⁵⁾ We expect that the difficulties can be overcome by introducing residual gauge fields depending only on x^0 . To study that possibility, we consider constructing an operator solution of the pure space-like axial gauge Schwinger model as a continuation from the light-cone gauge. We begin by constructing an operator solution in the region $0 \leq \theta < \frac{\pi}{4}$. This is straightforward since we can use the axial anomaly and the temporal gauge canonical quantization conditions as guiding principles in constructing the operator solution. We continue the operator solution into the axial region $\frac{\pi}{4} < \theta < \frac{\pi}{2}$. In the axial region the operator $(\partial_-)^{-1}$ becomes singular and gives rise to infrared divergences. Furthermore, in the axial region the Laplace operator, $m^2 - n_- \partial_+^2$ (where $m^2 = \frac{e^2}{\pi}$ and $n_- = \cos 2\theta$), becomes hyperbolic so that divergences result from its inversion. We find that the former divergences are canceled by the latter ones if we choose the representation of the residual gauge fields in such a way as to realize the ML prescription. As a consequence we recover well-defined fermion field operators. Bilinear products of these give rise to bosonized expressions for the vector current and the energy-momentum tensor through the equal x^+ -time point-splitting procedure. We apply McCartor and Robertson's method⁸⁾ and obtain the Hamiltonian for the residual gauge fields. It consists of a part from the physical operators integrated over $x^+ = \text{constant}$ and a part from the residual gauge operators integrated over $x^- = \text{constant}$. This is the way we extend the traditional Hamiltonian formulation to obtain the Hamiltonian for the residual gauge fields. Now it is straightforward to obtain an operator solution and the Hamiltonian in the gauge $A_1 = 0$ (here, the Coulomb gauge) in the ordinary coordinates. All one has to do is to take the

limit $\theta \rightarrow \frac{\pi}{2} - 0$. While that limit exists, we cannot simply set θ equal to $\frac{\pi}{2}$. For that reason one might say that we have not really quantized the Schwinger model at equal time in the Coulomb gauge. On the other hand one might regard θ as a regulating parameter. In that view we see that, as was the case in the light-cone gauge^{9,16)}, proper regulation of the theory necessarily involves information off the initial value surface. In any event, the solution contains the auxiliary fields and the representation space is of indefinite metric. This provides further indication that “physical” gauges are no more physical than “unphysical” gauges in that the representation space contains unphysical states and is of indefinite metric in all cases.

The paper is organized as follows: In § 2, we use the axial anomaly and the temporal gauge quantization conditions as guiding principles to construct the temporal gauge operator solution of the Schwinger model in the auxiliary coordinates. We also point out that if we choose the representation of the residual gauge fields so as to realize the PV prescription, we can construct fermion field operators in such a way that vacuum expectation values of them vanish. In § 3, we continue the solution into the axial region and show in detail that the infrared divergences inherent to the axial gauge quantizations are canceled if we choose the representation of the residual gauge fields which realizes the ML prescription. We also discuss properties of the axial gauge solution. Section 4 is devoted to concluding remarks.

We use the following conventions:

$$\begin{aligned}
g_{--} &= \cos 2\theta, & g_{-+} &= g_{+-} = \sin 2\theta, & g_{++} &= -\cos 2\theta, \\
g^{--} &= \cos 2\theta, & g^{-+} &= g^{+-} = \sin 2\theta, & g^{++} &= -\cos 2\theta, \\
n^\mu &= (n^+, n^-) = (0, 1), & n_\mu &= (n_+, n_-) = (\sin 2\theta, \cos 2\theta), \\
\gamma^0 &= \sigma_1, & \gamma^1 &= i\sigma_2, & \gamma^5 &= -\sigma_3, \\
\gamma^+ &= \gamma^0 \sin \theta + \gamma^1 \cos \theta = \begin{pmatrix} 0 & \sqrt{1+n_+} \\ -\varepsilon(n_-)\sqrt{1-n_+} & 0 \end{pmatrix}, \\
\gamma^- &= \gamma^0 \cos \theta - \gamma^1 \sin \theta = \begin{pmatrix} 0 & \varepsilon(n_-)\sqrt{1-n_+} \\ \sqrt{1+n_+} & 0 \end{pmatrix}.
\end{aligned}$$

§2. Temporal gauge formulation in the auxiliary coordinates

In this section we confine ourselves to the region $0 \leq \theta < \frac{\pi}{4}$ and choose x^- as the evolution parameter. The temporal gauge Schwinger model in the auxiliary coordinates is defined by the Lagrangian

$$L = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - B(n \cdot A) + i\bar{\psi}\gamma^\mu(\partial_\mu + ieA_\mu)\psi \quad (2.1)$$

where B is the Lagrange multiplier field, that is, the Nakanishi-Lautrup field in noncovariant formulations.¹⁷⁾ From the Lagrangian we derive the field equations

$$\partial_\mu F^{\mu\nu} = n^\nu B + J^\nu, \quad J^\nu = e\bar{\psi}\gamma^\nu\psi \quad (2.2)$$

$$i\gamma^\mu(\partial_\mu + ieA_\mu)\psi = 0, \quad (2.3)$$

and the gauge fixing condition

$$A_- = 0. \quad (2.4)$$

The field equation of B ,

$$\partial_- B = 0, \quad (2.5)$$

is obtained by operating on (2.2) with ∂_ν . Canonically conjugate momenta are defined by

$$\pi^- = \frac{\delta L}{\delta \partial_- A_-} = 0, \quad \pi^+ = \frac{\delta L}{\delta \partial_- A_+} = F_{-+}, \quad \pi_B = \frac{\delta L}{\delta \partial_+ B} = 0, \quad \pi_\psi = \frac{\delta L}{\delta \partial_- \psi} = i\bar{\psi}\gamma^-. \quad (2.6)$$

Thus we impose the following equal x^- -time canonical quantization conditions:

$$[A_+(x), A_+(y)] = [\pi^+(x), \pi^+(y)] = 0, \quad [A_+(x), \pi^+(y)] = i\delta(x^+ - y^+), \quad (2.7)$$

$$[A_+(x), \psi(y)] = [\pi^+(x), \psi(y)] = 0, \quad (2.8)$$

$$\{\psi_\alpha(x), \psi_\beta(y)\} = 0, \quad \{\psi_\alpha(x), \psi_\beta^*(y)\} = ((\gamma^0\gamma^-)^{-1})_{\alpha\beta}\delta(x^+ - y^+). \quad (2.9)$$

We begin constructing an operator solution by imposing the chiral anomaly

$$\partial_- J_+ - \partial_+ J_- = -m^2 F_{-+}. \quad (2.10)$$

The current operator given by (2.2) is

$$J_- = g_{-+}J^+ + g_{--}J^- = -\partial^+ F_{-+} - n_- B, \quad J_+ = \partial^- F_{-+} - n_+ B. \quad (2.11)$$

Substituting these expressions into (2.10) yields

$$(\partial_\mu \partial^\mu + m^2)F_{-+} = -n_- \partial_+ B. \quad (2.12)$$

Because $\partial_- B = 0$, F_{-+} is solved to be

$$F_{-+} = \partial_- A_+ = -m\tilde{\Sigma} - \frac{n_-}{m^2 - n_- \partial_+^2} \partial_+ B. \quad (2.13)$$

Here we have identified the homogeneous solution with the massive free field $\tilde{\Sigma}$, since F_{-+} is gauge invariant. The normalization is determined in such a way that it agrees with the Landau gauge solution¹⁸⁾. $\tilde{\Sigma}$ is given by

$$\tilde{\Sigma}(x) = \frac{1}{\sqrt{4\pi}} \int_{-\infty}^{\infty} \frac{dp_+}{\sqrt{p^-}} \{a(p_+)e^{-ip \cdot x} + a^*(p_+)e^{ip \cdot x}\}, \quad (2.14)$$

where $p^- = \sqrt{p_+^2 + m_0^2}$ with $m_0^2 = n_- m^2$ and

$$[a(p_+), a(q_+)] = 0, \quad [a(p_+), a^*(q_+)] = \delta(p_+ - q_+). \quad (2.15)$$

By integrating (2.13) with respect to x^- we obtain the general solution for A_+ as

$$A_+ = -\frac{m}{\partial_-} \tilde{\Sigma} - \frac{n_- x^- \partial_+}{m^2 - n_- \partial_+^2} B + \text{integration constant}. \quad (2.16)$$

We determine the integration constant in the following way: From the form of the d'Alembertian in the auxiliary coordinates,

$$\partial_\mu \partial^\mu = n_- \partial_-^2 + 2n_+ n_- \partial_+ \partial_- - n_- \partial_+^2 \quad (2.17)$$

we rewrite the first term of (2.16) as

$$\frac{1}{\partial_-} \tilde{\Sigma} = -\frac{n_+ \partial_+ + \partial_-}{m^2 - n_- \partial_+^2} \tilde{\Sigma}. \quad (2.18)$$

In the range of θ we are considering, we see that the operator $(\partial_-)^{-1}$ does not give rise to any infrared divergences. Now the first term in (2.16) has the property $[\frac{m}{\partial_-} \tilde{\Sigma}(x^+), \frac{m}{\partial_-} \tilde{\Sigma}(y^+)] \neq 0$ at equal x^- -time because the $\tilde{\Sigma}$ satisfies the following equal x^- -time commutation relations

$$[\tilde{\Sigma}(x), \tilde{\Sigma}(y)] = [\partial^- \tilde{\Sigma}(x), \partial^- \tilde{\Sigma}(y)] = 0, \quad [\tilde{\Sigma}(x), \partial^- \tilde{\Sigma}(y)] = i\delta(x^+ - y^+). \quad (2.19)$$

Therefore, to preserve the canonical commutation relations (2.7), we must cancel the nonvanishing terms by contributions from another field C , the integration constant, which depends only on x^+ . We find that if B and C satisfy the commutation relations

$$\begin{aligned} [B(x), B(y)] &= [C(x), C(y)] = 0, \quad [B(x), C(y)] = i(m^2 - n_- \partial_+^2) \delta(x^+ - y^+), \\ [B(x), \tilde{\Sigma}(y)] &= [C(x), \tilde{\Sigma}(y)] = 0, \end{aligned} \quad (2.20)$$

then

$$A_+ = -\frac{m}{\partial_-} \tilde{\Sigma} + \frac{\partial_+}{m^2 - n_- \partial_+^2} (C - n_- x^- B) + \frac{n_+ m^2}{(m^2 - n_- \partial_+^2)^2} B \quad (2.21)$$

satisfies the quantization conditions in (2.7). In this way we can determine the integration constant: by taking account of the canonical quantization conditions in the temporal gauge formulation.

Now that we have the A_+ , we proceed to solve the fermion field equation (2.3). A formal solution is given by

$$\psi_\alpha(x) = \frac{Z_\alpha}{\sqrt{(\gamma^0 \gamma^-)_{\alpha\alpha}}} \exp[-i\sqrt{\pi} \Lambda_\alpha(x)], \quad (2.22)$$

where Z_α is normalization constant and

$$A_\alpha(x) = X(x) + (-1)^\alpha \lambda(x) \quad (2.23)$$

with

$$\begin{aligned} X &= \frac{\partial^+}{\partial_-} \tilde{\Sigma} + \frac{m}{m^2 - n_- \partial_+^2} \left(C - n_- x^- B + \frac{n_+ n_-}{m^2 - n_- \partial_+^2} \partial_+ B \right), \\ \lambda &= \tilde{\Sigma} + \frac{m}{m^2 - n_- \partial_+^2} \partial_+^{-1} B. \end{aligned} \quad (2.24)$$

Here the operator ∂_+^{-1} is defined by

$$\partial_+^{-1} B(x) = \frac{1}{2} \int_{-\infty}^{\infty} dy^+ \varepsilon(x^+ - y^+) B(y), \quad (2.25)$$

which imposes, in effect, the principal value regularization. To further examine the properties of the fermion field operators, such as their vacuum expectation values, we need explicit representations of the B and C . Note that the canonical commutation relations cannot uniquely determine the representations of the canonical variables, as was pointed out by Haller¹⁹⁾. One possibility is to choose the following zero norm and positive metric expressions for B and C :

$$\begin{aligned} B(x) &= \frac{m}{\sqrt{4\pi}} \int_{-\infty}^{\infty} dk_+ \sqrt{|k_+|} \{ (a_0(k_+) - a_1(k_+)) e^{-ik_+ x^+} + \text{h.c.} \}, \\ \frac{m}{m^2 - n_- \partial_+^2} C(x) &= \frac{i}{\sqrt{4\pi}} \int_{-\infty}^{\infty} \frac{dk_+}{\sqrt{|k_+|}} \{ a_1(k_+) e^{-ik_+ x^+} - a_1^*(k_+) e^{ik_+ x^+} \}, \end{aligned} \quad (2.26)$$

where

$$[a_0(k_+), a_0^*(q_+)] = -\delta(k_+ - q_+), \quad [a_1(k_+), a_1^*(q_+)] = \delta(k_+ - q_+), \quad (2.27)$$

and all other commutators are zero. It can be shown that B and C in (2.26) realize the PV prescription. With this choice, the spurion operators are composed out of ultraviolet parts of the $\tilde{\Sigma}$ as well as infrared parts of C and $\partial_+^{-1} B$ according to the method given in Ref.18). This in turn gives rise to vanishing vacuum expectation values for the fermion field operators. In the Landau gauge solution¹⁸⁾ the fermion fields have vanishing vacuum expectation value but in the light-cone gauge solution the vacuum expectation value of ψ_+ diverges. The inclusion of the ultraviolet parts of $\tilde{\Sigma}$ in the spurions causes them to become x dependent and the solution has some different characteristics from the previous solutions. We are not completely certain whether the solution using the choice (2.26) is completely consistent but we do know that when we move to the axial gauge choice ($\theta > \frac{\pi}{4}$) we cannot

use the representation (2.26): if we do there are uncanceled infrared divergences which destroy the solution. We will not further discuss the choice (2.26) in the present paper but will concentrate on another choice, one which realizes the ML prescription. To do that we must limit the k_+ -integration region to be $(0, \infty)$; we showed in Ref.13) that by choosing a suitable vacuum it is always possible to do so. We show in the next section that with the new choice, the infrared divergences in the axial gauge cancel.

To implement the new(ML) representation we choose the following representation for B and C :

$$\begin{aligned} B(x) &= \frac{m}{\sqrt{2\pi}} \int_0^\infty dk_+ \sqrt{k_+} \{ B(k_+) e^{-ik_+ x^+} + B^*(k_+) e^{ik_+ x^+} \}, \\ \frac{m}{m^2 - n_- \partial_+^2} C(x) &= \frac{i}{\sqrt{2\pi}} \int_0^\infty \frac{dk_+}{\sqrt{k_+}} \{ C(k_+) e^{-ik_+ x^+} - C^*(k_+) e^{ik_+ x^+} \}, \end{aligned} \quad (2.28)$$

where

$$[B(k_+), C^*(q_+)] = [C(k_+), B^*(q_+)] = -\delta(k_+ - q_+), \quad (2.29)$$

and all other commutators are zero. Note that the C becomes a zero norm field as a consequence of altering the representations. Note also that the vacuum expectation value of ψ_2 diverges as it does in the light-cone gauge solution⁹⁾.

We define the physical subspace, V , by

$$V = \{ |\text{phys}\rangle \mid B(k_+) |\text{phys}\rangle = 0 \}. \quad (2.30)$$

To subtract the infrared parts of the massless fields into the spurion operators σ_α , we define the infrared parts, $\Lambda_\alpha^{(0)}$, of Λ_α to be

$$\Lambda_\alpha^{(0)} = \frac{i}{\sqrt{2\pi}} \int_0^\kappa \frac{dk_+}{\sqrt{k_+}} \{ C(k_+) - C^*(k_+) + (-1)^\alpha (B(k_+) - B^*(k_+)) \} \quad (2.31)$$

where κ is a small positive constant. Then the fermion field operators are defined as

$$\psi_\alpha(x) = \frac{Z_\alpha}{\sqrt{(\gamma^0 \gamma^-)_{\alpha\alpha}}} \exp[-i\sqrt{\pi} \Lambda_{\alpha r}^{(-)}(x)] \sigma_\alpha \exp[-i\sqrt{\pi} \Lambda_{\alpha r}^{(+)}(x)] \quad (2.32)$$

where $\Lambda_{\alpha r}^{(-)}$ and $\Lambda_{\alpha r}^{(+)}$ are the creation and annihilation operator parts of $\Lambda_{\alpha r} \equiv \Lambda_\alpha - \Lambda_\alpha^{(0)}$ and

$$\sigma_\alpha = \exp \left[-i\sqrt{\pi} \left(\Lambda_\alpha^{(0)} - (-1)^\alpha \frac{Q}{2m} \right) \right] \quad (2.33)$$

Here, $Q = -\int_{-\infty}^\infty dx^+ B(x)$; note that Q in σ_α constitutes a Klein transformation.

We enumerate properties of the ψ_α to show that the ψ_α constitute the operator solution of the temporal gauge Schwinger model in the auxiliary coordinates.

(1) The Dirac equation is satisfied:

$$i\gamma^\mu(\partial_\mu + ieA_\mu)\psi = 0 \quad (2.34)$$

To get this we use the fact that $[A_+^{(+)}(x), \psi_\alpha(x)] = 0$, where $A_+^{(+)}$ is the annihilation operator part of A_+ .

(2) The canonical commutation relations (2.8) and anticommutation relations (2.9) are satisfied.

(3) By applying the gauge invariant point-splitting procedure to $e\bar{\psi}\gamma^\mu\psi$, we obtain the vector current $J^\mu = m\varepsilon^{\mu\nu}\partial_\nu\lambda$, where $\varepsilon^{++} = \varepsilon^{--} = 0$, $\varepsilon^{+-} = -\varepsilon^{-+} = 1$. The charge operator, Q , is given by

$$Q = \int_{-\infty}^{\infty} dx^+ J^-(x) = - \int_{-\infty}^{\infty} dx^+ B(x), \quad (2.35)$$

where the derivative terms integrate to zero.

(4) The conserved chiral current $J^{\mu 5} = m\varepsilon^{\mu\nu}\partial_\nu X$ is similarly obtained from $e\bar{\psi}\gamma^\mu\gamma^5\psi$. The chiral charge operator, Q_5 , is given by

$$Q_5 = - \int_{-\infty}^{\infty} dx^+ J^{-5}(x) = \int_{-\infty}^{\infty} dx^+ \partial_+ C(x), \quad (2.36)$$

where the derivative terms, except $\partial_+ C$, integrate to zero. The term $\partial_+ C$ does not integrate to zero because ψ contains the singular operator $\partial_+^{-1}B$.

(5) Applying the gauge invariant point-splitting procedure to the fermi products in the symmetric energy-momentum tensor and subtracting a diverging c-number, which we denote by R , we get

$$\begin{aligned} \Theta_+^- &= \frac{i}{2}R(\bar{\psi}\gamma^- \partial_+ \psi - \partial_+ \bar{\psi}\gamma^- \psi) - A_+ J^- - \frac{1}{2}(A_+, B)_+ \\ &= \frac{1}{2}(\partial_+ \lambda, \partial_- \lambda)_+ - \frac{1}{2}(A_+, B)_+, \end{aligned} \quad (2.37)$$

$$\begin{aligned} \Theta_+^+ &= \frac{i}{2}R(\bar{\psi}\gamma^+ \partial_+ \psi - \partial_+ \bar{\psi}\gamma^+ \psi) - A_+ J^+ + \frac{1}{2}(F_{-+})^2 \\ &= -\frac{n_-}{2}\{(\partial_+ \lambda)^2 + (\partial_- \lambda)^2\} + \frac{1}{2}(F_{-+})^2, \end{aligned} \quad (2.38)$$

$$\begin{aligned} \Theta_-^- &= -\frac{i}{2}R(\bar{\psi}\gamma^+ \partial_+ \psi - \partial_+ \bar{\psi}\gamma^+ \psi) + A_+ J^+ + \frac{1}{2}(F_{-+})^2 \\ &= \frac{n_-}{2}\{(\partial_+ \lambda)^2 + (\partial_- \lambda)^2\} + \frac{1}{2}(F_{-+})^2, \end{aligned} \quad (2.39)$$

$$\begin{aligned}
\Theta_-^+ &= -\frac{in_+}{2n_-}R(\bar{\psi}\gamma^+\partial_+\psi - \partial_+\bar{\psi}\gamma^+\psi) - \frac{i}{2n_-}R(\bar{\psi}\gamma^+\gamma^5\partial_+\psi - \partial_+\bar{\psi}\gamma^+\gamma^5\psi) \\
&+ \frac{1}{n_-}J_-A_+ + \frac{n_+}{n_-}J^+A_+ = \frac{1}{2}(\partial_-\lambda, \partial^+\lambda)_+.
\end{aligned} \tag{2.40}$$

(6) Translational generators consist of those of the constituent fields:

$$\begin{aligned}
P_- &= \int_{-\infty}^{\infty} dx^+ : \Theta_-^- : \\
&= \int_{-\infty}^{\infty} dx^+ \frac{1}{2} : \{n_-(\partial_+\tilde{\Sigma})^2 + n_-(\partial_-\tilde{\Sigma})^2 + m^2(\tilde{\Sigma})^2 + B\frac{n_-}{m^2 - n_-\partial_+^2}B\} :, \\
P_+ &= \int_{-\infty}^{\infty} dx^+ : \Theta_+^- := \int_{-\infty}^{\infty} dx^+ : \{\partial_+\tilde{\Sigma}\partial^-\tilde{\Sigma} - B\frac{1}{m^2 - n_-\partial_+^2}\partial_+C\} : .
\end{aligned} \tag{2.41}$$

§3. Cancellation of Infrared divergences resulting from ∂_-^{-1}

In this section we continue θ into the region $\frac{\pi}{4} < \theta < \frac{\pi}{2}$ and take x^+ as the evolution parameter. In accordance with the change of the evolution parameter we change the Fourier expansion of $\tilde{\Sigma}$ so as to be convenient for calculating equal x^+ -time commutation relations. By making use of $\frac{dp_+}{p^-} = -\frac{dp_-}{p^+}$ we change integration variables from p_+ to p_- and rewrite $\tilde{\Sigma}$ as follows

$$\tilde{\Sigma}(x) = \frac{1}{\sqrt{4\pi}} \int_{-\infty}^{\infty} \frac{dp_-}{\sqrt{p^+}} \{a(p_-)e^{-ip \cdot x} + a^*(p_-)e^{ip \cdot x}\}, \tag{3.1}$$

where $p^+ = \sqrt{p_-^2 + m_0^2}$ with $m_0^2 = |n_-| m^2$ and

$$[a(p_-), a(q_-)] = 0, \quad [a(p_-), a^*(q_-)] = \delta(p_- - q_-). \tag{3.2}$$

$\tilde{\Sigma}$ satisfies the following equal x^+ -time commutation relations:

$$[\tilde{\Sigma}(x), \tilde{\Sigma}(y)] = [\partial^+\tilde{\Sigma}(x), \partial^+\tilde{\Sigma}(y)] = 0, \quad [\tilde{\Sigma}(x), \partial^+\tilde{\Sigma}(y)] = i\delta(x^- - y^-). \tag{3.3}$$

Furthermore, in accordance with the change of the evolution parameter, the conjugate momentum of ψ changes to

$$\pi_\psi^{(a)} = \frac{\delta L}{\delta \partial_+\psi} = i\bar{\psi}\gamma^+. \tag{3.4}$$

Therefore, we have to modify the normalization of ψ in (2.32) accordingly:

$$\psi_\alpha^{(a)}(x) = \frac{Z_\alpha^{(a)}}{\sqrt{(\gamma^0\gamma^+)_{\alpha\alpha}}} \exp[-i\sqrt{\pi}\Lambda_{\alpha r}^{(-)}(x)]\sigma_\alpha \exp[-i\sqrt{\pi}\Lambda_{\alpha r}^{(+)}(x)], \tag{3.5}$$

where we have put (a) on ψ and on the normalization constant to denote that the normalization is altered.

We begin by pointing out that because $n_- = \cos 2\theta < 0$ in the region $\frac{\pi}{4} < \theta < \frac{\pi}{2}$, the Laplace operator, $m^2 - n_- \partial_+^2$, which operates on the residual gauge fields, becomes hyperbolic so that its inverse gives rise to divergences. Therefore, we regularize them by PV regularization. We then recover well-defined fermion field operators because linear infrared divergences resulting from $\frac{\partial_\pm}{\partial_-} \tilde{\Sigma}$ are canceled by those resulting from the square of the inverse of the hyperbolic Laplace operator. To show that this really happens, we rewrite $\psi_\alpha^{(a)*}(x)\psi_\alpha^{(a)}(y)$ into normal ordered form. It is tedious but straightforward to obtain

$$\begin{aligned} \psi_\alpha^{(a)*}(x)\psi_\alpha^{(a)}(y) &= \frac{(Z_\alpha^{(a)})^2}{(\gamma^0 \gamma^+)_{\alpha\alpha}} \exp[M_{\alpha\alpha}(x-y)] \exp[-i\sqrt{\pi}(\Lambda_\alpha^{(-)}(x) - \Lambda_\alpha^{(-)}(y))] \\ &\quad \times \exp[-i\sqrt{\pi}(\Lambda_\alpha^{(+)}(x) - \Lambda_\alpha^{(+)}(y))], \end{aligned} \quad (3.6)$$

where

$$\begin{aligned} M_{\alpha\alpha}(x) &= \frac{1}{4} \int_{-\infty}^{\infty} \frac{dp_-}{p^+} \left(1 + \left(\frac{p^+}{p_-} \right)^2 + 2(-1)^\alpha \frac{p^+}{p_-} \right) e^{-ip \cdot x} \\ &\quad + \frac{1}{2} \int_0^{\infty} dk_+ \left(-\frac{im^2 n_- x^-}{m^2 + n_- k_+^2} + \frac{2m^2 n_- n_+ k_+}{(m^2 + n_- k_+^2)^2} \right) e^{-ik_+ x^+} \\ &\quad + (-1)^\alpha \int_0^{\infty} dk_+ \left(\frac{n_- k_+}{m^2 + n_- k_+^2} e^{-ik_+ x^+} - \frac{e^{-ik_+ x^+} - \theta(\kappa - k_+)}{k_+} \right). \end{aligned} \quad (3.7)$$

We must show that $M_{\alpha\alpha}(x)$ does not diverge when $x^+ = 0$, because in what follows we make use of the values of $M_{\alpha\alpha}(x-y)$ at $x^+ = y^+$ to calculate the equal x^+ -time anticommutation relations and to calculate the dynamical operators.

Since $(p^+)^2 = p_-^2 + m_0^2$, only the term $\frac{m_0^2}{p_-^2}$ among the physical contributions on the first line of (3.7) gives rise to a linear infrared divergence. We rewrite the expression as follows

$$\frac{1}{4} \int_{-\infty}^{\infty} \frac{dp_-}{p^+} \frac{m_0^2}{p_-^2} e^{-ip_- x^-} = \int_0^{\infty} \frac{dq}{q^2} - \frac{m_0^2}{2} \int_0^{\infty} \frac{dp_-}{p^+} \frac{1 - \cos p_- x^-}{p_-^2}, \quad (3.8)$$

where the first, linearly diverging integration is obtained by changing integration variables from p_- to q through $p_- = \frac{m_0 q}{2\sqrt{q+1}}$.

As for the residual gauge field's contributions: those divergences which can be regularized by the PV prescription we regularize by the PV prescription. Then we find that when $x^+ = 0$, the first term on the second line of (3.7) is equal to zero:

$$\int_0^{\infty} dk_+ \frac{n_- m^2}{m^2 + n_- k_+^2} = \int_0^{\infty} dk_+ \frac{m^2}{k_+^2 - a^2} = 0, \quad (3.9)$$

where $a = \frac{m}{\sqrt{-n_-}}$. However, we cannot regularize a linear divergence resulting from a double pole by the PV regularization. Therefore when $x^+ = 0$, the second term on the second line

of (3.7) gives rise to a linear divergence as follows

$$\begin{aligned} \int_0^\infty dk_+ \frac{n_- n_+ m^2 k_+}{(m^2 + n_- k_+^2)^2} &= -\frac{n_+ a}{4} \int_0^\infty dk_+ \left(\frac{1}{(k_+ - a)^2} - \frac{1}{(k_+ + a)^2} \right) \\ &= -\frac{n_+ a}{4} \left(\int_0^a dk \frac{1}{k^2} + \int_0^\infty dk \frac{1}{k^2} - \int_a^\infty dk \frac{1}{k^2} \right) = -\int_0^\infty dq \frac{1}{q^2} + \frac{n_+}{2}, \end{aligned} \quad (3.10)$$

where the last, linearly diverging term is given by changing integration variables from k to q through $k = \frac{n_+ a}{2} q$. Thus we see that the linear divergence originated from the physical contributions is canceled by that resulting from the residual gauge field's contributions. It should be noted here that if we had chosen the representation of the residual gauge fields so as to realize the PV prescription (the expression given by (2.26)), we would obtain the identically vanishing integral $\int_{-\infty}^\infty dk_+ \frac{n_- n_+ m^2 k_+}{(m^2 + n_- k_+^2)^2} = 0$ so that the linear divergence resulting from $\frac{\partial^+}{\partial_-} \tilde{\Sigma}$ is not canceled.

It turns out that when $x^+ = 0$, the terms on the third line of (3.7) give rise to a finite value, $\log \frac{\kappa}{\sqrt{|\kappa^2 - a^2|}}$, so that by summing all the finite contributions we obtain

$$\begin{aligned} M_{\alpha\alpha}(x)|_{x^+=0} &= K_0(m_0|x^-|) + i(-1)^{\alpha-1} \frac{\pi}{2} \varepsilon(x^-) \\ &\quad - \frac{m_0^2}{2} \int_0^\infty \frac{dp_-}{p^+} \frac{1 - \cos p_- x^-}{p_-^2} + \frac{n_+}{2} + (-1)^\alpha \log \frac{\kappa}{\sqrt{|\kappa^2 - a^2|}} \\ &= -\left(\log \frac{m_0 \gamma}{2} + \log(x^- + i\varepsilon(-1)^{\alpha-1}) \right) + i(-1)^{\alpha-1} \frac{\pi}{2} + F_\alpha(|x^-|), \end{aligned} \quad (3.11)$$

where $K_0(m_0|x^-|)$ is the modified Bessel function of order 0, which possesses a logarithmic term: $-(\log \frac{m_0|x^-|}{2} + \gamma)$. We have gathered the remainder of K_0 and the other terms to make a continuous function $F_\alpha(|x^-|)$.

Now that we have finished showing that the linear divergences cancel each other, we proceed to study other properties of the fermion field operators.

(1) The equal x^+ -time anticommutation relations hold. For example, when $x^+ = y^+$, we find that

$$\exp[M_{\alpha\alpha}(x - y)] = \frac{2}{m_0 e^\gamma} \frac{i(-1)^{\alpha-1}}{x^- - y^- + i\varepsilon(-1)^{\alpha-1}} e^{F_\alpha(|x^- - y^-|)}, \quad (3.12)$$

so that if we take $Z_\alpha^{(a)}$ to be

$$(Z_\alpha^{(a)})^2 = \frac{m_0 e^\gamma}{4\pi} e^{-F_\alpha(0)}, \quad (3.13)$$

then we obtain

$$\begin{aligned} &\{\psi_\alpha^*(x), \psi_\alpha(y)\}|_{x^+=y^+} \\ &= \frac{\exp[F_\alpha(|x^- - y^-|) - F_\alpha(0)]}{2\pi(\gamma^0 \gamma^+)_{\alpha\alpha}} \left(\frac{i(-1)^{\alpha-1}}{x^- - y^- + i\varepsilon(-1)^{\alpha-1}} + \frac{i(-1)^{\alpha-1}}{y^- - x^- + i\varepsilon(-1)^{\alpha-1}} \right) \end{aligned}$$

$$\begin{aligned}
& \times \exp[-i\sqrt{\pi}(\Lambda_{\alpha}^{(-)}(x) - \Lambda_{\alpha}^{(-)}(y))]\exp[-i\sqrt{\pi}(\Lambda_{\alpha}^{(+)}(x) - \Lambda_{\alpha}^{(+)}(y))] \\
& = \frac{1}{(\gamma^0\gamma^+)_{\alpha\alpha}}\delta(x^- - y^-).
\end{aligned} \tag{3.14}$$

When $\alpha \neq \beta$, $\{\psi_{\alpha}^*(x), \psi_{\beta}(y)\}|_{x^+=y^+}$ vanishes due to the Klein transformation factors.

(2) The vector current, $J^{\mu} = m\varepsilon^{\mu\nu}\partial_{\nu}\lambda$, is obtained this time by applying the equal x^+ -time point-splitting procedure to $e\bar{\psi}\gamma^{\mu}\psi$.

(3) The conserved chiral current $J^{\mu 5} = m\varepsilon^{\mu\nu}\partial_{\nu}X$ is similarly obtained from $e\bar{\psi}\gamma^{\mu}\gamma^5\psi$.

(4) As a consequence of our choice of representation of the residual gauge fields, we obtain the ML form of the x^+ -ordered gauge field propagator:

$$\begin{aligned}
D_{\mu\nu}(x-y) &= \langle 0 | \{ \theta(x^+ - y^+) A_{\mu}(x) A_{\nu}(y) + \theta(y^+ - x^+) A_{\nu}(y) A_{\mu}(x) \} | 0 \rangle \\
&= \frac{1}{(2\pi)^2} \int d^2q D_{\mu\nu}(q) e^{-iq \cdot (x-y)}
\end{aligned} \tag{3.15}$$

where

$$\begin{aligned}
D_{\mu\nu}(q) &= \frac{i}{q^2 - m^2 + i\epsilon} \left(-g_{\mu\nu} + \frac{n_{\mu}q_{\nu} + n_{\nu}q_{\mu}}{q_- + i\varepsilon \text{sgn}(q_+)} - n^2 \frac{q_{\mu}q_{\nu}}{(q_- + i\varepsilon \text{sgn}(q_+))^2} \right) \\
&\quad - \delta_{\mu+}\delta_{\nu+} \frac{i}{2} \left(\frac{1}{(q_- + i\varepsilon)^2} + \frac{1}{(q_- + i\varepsilon)^2} \right).
\end{aligned} \tag{3.16}$$

We give a detailed derivation of this in the Appendix.

(5) Applying the gauge invariant point-splitting procedure to the fermi products in the symmetric energy-momentum tensor and subtracting a diverging c-number, which we denote by R , we get

$$\begin{aligned}
\Theta_+^- &= \frac{in_+}{2n_-} R(\bar{\psi}\gamma^- \partial_- \psi - \partial_- \bar{\psi}\gamma^- \psi) - \frac{i}{2n_-} R(\bar{\psi}\gamma^- \gamma^5 \partial_- \psi - \partial_- \bar{\psi}\gamma^- \gamma^5 \psi) \\
&\quad - \frac{1}{2}(A_+, B)_+ = \frac{1}{2}(\partial_+ \lambda, \partial^- \lambda)_+ - \frac{1}{2}(A_+, B)_+,
\end{aligned} \tag{3.17}$$

$$\begin{aligned}
\Theta_+^+ &= -\frac{i}{2} R(\bar{\psi}\gamma^- \partial_- \psi - \partial_- \bar{\psi}\gamma^- \psi) + \frac{1}{2}(F_{-+})^2 \\
&= -\frac{n_-}{2} \{(\partial_+ \lambda)^2 + (\partial_- \lambda)^2\} + \frac{1}{2}(F_{-+})^2,
\end{aligned} \tag{3.18}$$

$$\begin{aligned}
\Theta_-^- &= \frac{1}{2}(F_{-+})^2 + \frac{i}{4} R(\bar{\psi}\gamma^- \partial_- \psi - \partial_- \bar{\psi}\gamma^- \psi) \\
&\quad - \frac{n_+}{4} R(\bar{\psi}\gamma^- \gamma^5 \partial_- \psi - \partial_- \bar{\psi}\gamma^- \gamma^5 \psi) + \frac{n_-}{4} R(\bar{\psi}\gamma^+ \gamma^5 \partial_- \psi - \partial_- \bar{\psi}\gamma^+ \gamma^5 \psi) \\
&= \frac{n_-}{2} \{(\partial_+ \lambda)^2 + (\partial_- \lambda)^2\} + \frac{1}{2}(F_{-+})^2,
\end{aligned} \tag{3.19}$$

$$\Theta_-^+ = \frac{i}{2} R(\bar{\psi}\gamma^+\partial_-\psi - \partial_-\bar{\psi}\gamma^+\psi) = \frac{1}{2}(\partial_-\lambda, \partial^+\lambda)_+. \quad (3.20)$$

We must take special care in the axial gauge formulation when we derive the conserved translational generators from the divergence equation

$$\partial_\nu \Theta_\mu{}^\nu = 0. \quad (3.21)$$

The problem is that although x^- is a space coordinate, Θ_μ^- does not vanish in the limits $x^- \rightarrow \pm\infty$. This is because the residual gauge fields do not depend on x^- and because A_+ depends explicitly on x^- . Therefore we have to retain $\lim_{x^- \rightarrow \pm\infty} \Theta_\mu^-$. To take this fact into account, we integrate the divergence equation $\partial_\nu \Theta_\mu{}^\nu = 0$ over a closed surface shown in Fig. 1, whose bounds T and L are let tend to ∞ after all calculations are finished. We remark that we can use this surface even in the limit $\theta \rightarrow \frac{\pi}{2} - 0$, in contrast with one used in Refs. 8) and 20).




Fig. 1.

It is straightforward to obtain

$$0 = \left(\int_{-L}^L dx^- [\Theta_\mu^+(x)]_{x^+=-T}^{x^+=T} + \int_{-T}^T dx^+ [\Theta_\mu^-(x)]_{x^-=-L}^{x^-=L} \right), \quad (3.22)$$

where

$$\begin{aligned}\left[\Theta_{\mu}^{+}(x)\right]_{x^{+}=-T}^{x^{+}=T} &= \Theta_{\mu}^{+}(x)|_{x^{+}=T} - \Theta_{\mu}^{+}(x)|_{x^{+}=-T}, \\ \left[\Theta_{\mu}^{-}(x)\right]_{x^{-}=-L}^{x^{-}=L} &= \Theta_{\mu}^{-}(x)|_{x^{-}=L} - \Theta_{\mu}^{-}(x)|_{x^{-}=-L}.\end{aligned}\tag{3.23}$$

In what follows we refer to operators made of $\tilde{\Sigma}$ as physical operators and operators made of B and C as residual gauge operators.

It can be shown that products of physical operators and residual gauge operators vanish, so that the physical operator parts and the residual gauge operator parts decouple in (3.22). This result reflects the fact that the physical operator parts are conserved by themselves. It can be also shown that the residual gauge operators which are not well-defined in the limit $L \rightarrow \infty$ cancel among themselves. After getting the decoupled and well-defined expressions, we take the limit $L \rightarrow \infty$. This limit enables us to discard the physical operator parts $\theta_{\mu}^{-}(x)$ of $\Theta_{\mu}^{-}(x)$, as is usually done when one calculates the conserved physical generators in the traditional axial gauge formulations. Finally, we take the limit $T \rightarrow \infty$. At this stage we assume that we can integrate the remaining residual gauge operators in $\Theta_{\mu}^{-}(x)$ by parts in the x^{+} direction. This assumption is justified because the residual gauge operator parts are conserved by themselves. As a consequence, we obtain

$$0 = \int_{-\infty}^{\infty} dx^{-} \left[\theta_{+}^{+}\right]_{x^{+}=-\infty}^{x^{+}=\infty} - \int_{-\infty}^{\infty} dx^{+} \left[B \frac{1}{m^2 - n_{-} \partial_{+}^2} \partial_{+} C\right]_{x^{-}=-\infty}^{x^{-}=\infty}, \tag{3.24}$$

$$0 = \int_{-\infty}^{\infty} dx^{-} \left[\theta_{-}^{+}\right]_{x^{+}=-\infty}^{x^{+}=\infty} + \frac{1}{2} \int_{-\infty}^{\infty} dx^{+} \left[B(x) \frac{n_{-}}{m^2 - n_{-} \partial_{+}^2} B(x)\right]_{x^{-}=-\infty}^{x^{-}=\infty}, \tag{3.25}$$

where

$$\begin{aligned}\theta_{+}^{+} &= \frac{n_{+}}{n_{-}} : \partial_{-} \tilde{\Sigma} \partial^{+} \tilde{\Sigma} : - \frac{1}{2n_{-}} : \{(\partial^{+} \tilde{\Sigma})^2 + (\partial_{-} \tilde{\Sigma})^2 + m_0^2 (\tilde{\Sigma})^2\} :, \\ \theta_{-}^{+} &= : \partial_{-} \tilde{\Sigma} \partial^{+} \tilde{\Sigma} : .\end{aligned}\tag{3.26}$$

From these relations we obtain the conserved generators:

$$P_{+} = \int_{-\infty}^{\infty} dx^{-} \theta_{+}^{+}(x) - \int_{-\infty}^{\infty} dx^{+} B(x) \frac{1}{m^2 - n_{-} \partial_{+}^2} \partial_{+} C(x), \tag{3.27}$$

$$P_{-} = \int_{-\infty}^{\infty} dx^{-} \theta_{-}^{+}(x) + \frac{1}{2} \int_{-\infty}^{\infty} dx^{+} B(x) \frac{n_{-}}{m^2 - n_{-} \partial_{+}^2} B(x). \tag{3.28}$$

In this way, even in the axial gauge formulation, we can obtain the conserved translational generators. These consist of the physical operator part integrated over $x^{+} = \text{constant}$ and

the residual gauge operator part integrated over $x^- = \text{constant}$. By making use of the commutation relations (2.20) and (3.3), it is easy to show that the Heisenberg equations for A_+ and $\psi^{(a)}$ hold. This shows that we have successfully constructed the extended Hamiltonian formulation of the axial gauge Schwinger model.

Finally, we shall discuss the charges in the axial gauge formulation. The current, being gauge invariant, is the same as in the temporal gauge

$$J^- = -\partial_+(m\tilde{\Sigma} + \frac{m^2}{m^2 - n_- \partial_+^2} \partial_+^{-1} B), \quad J^+ = \partial_-(m\tilde{\Sigma} + \frac{m^2}{m^2 - n_- \partial_+^2} \partial_+^{-1} B) = m\partial_- \tilde{\Sigma}. \quad (3.29)$$

Using the fact that the current has zero divergence and also using the relation

$$\frac{m^2}{m^2 - n_- \partial_+^2} B = B + \frac{n_- \partial_+^2}{m^2 - n_- \partial_+^2} B, \quad (3.30)$$

we find the charge to be

$$Q = -\int_{-\infty}^{\infty} dx^+ B(x) + \int_{-\infty}^{\infty} dx^- m\partial_- \tilde{\Sigma}(x). \quad (3.31)$$

In the temporal gauge formulation (and also in the light-cone gauge⁹⁾ and Landau gauge¹⁸⁾ solutions) we set the second term equal to zero. The justification for that is that $\tilde{\Sigma}(x)$ is a massive field and since it has no infrared singularities, the term involving the integral over it commutes with the ψ field. The question is somewhat more complicated in the axial gauge case since the singularity coming from ∂_-^{-1} causes the commutator of the second term in (3.31) with fermi field to become nonzero. On the other hand, we cannot keep both terms in (3.31) since in that case the fermi field would carry twice the correct charge. We believe that the correct charge operator is obtained from (3.31) by letting the second term provide vanishing topological charges. In previous attempts to formulate the Coulomb gauge Schwinger model (using a representation space which is entirely physical), the fermion field operators were represented as solitons made solely out of the $\tilde{\Sigma}$ so that the $\tilde{\Sigma}$ tended to nonvanishing values in the limits $x^- \rightarrow \pm\infty$. That led, inevitably, to infrared difficulties. In our formulation, the fermion field operators are constituted in the same manner as ones in the temporal gauge formulation so that it is reasonable for them not to carry any topological charges. To attain this we define the operator $(\partial_-)^{-1}$ as a finite integral:

$$\frac{1}{2} \int_{-L}^L \varepsilon(x^- - y^-) f(y^-) dy^- \quad (3.32)$$

and take the limit $L \rightarrow \infty$ after all relevant calculations are finished. This procedure is justified by the fact that the infrared divergences are all eliminated. We therefore take the charge operator to be

$$Q = -\int_{-\infty}^{\infty} dx^+ B(x). \quad (3.33)$$

Similarly, we take the chiral charge to be

$$Q_5 = \int_{-\infty}^{\infty} dx^+ \partial_+ C(x). \quad (3.34)$$

§4. Concluding remarks

In this paper we have shown that x^- -independent residual gauge fields can be introduced as regulator fields in the pure space-like axial gauge formulation of the Schwinger model by extending the Hamiltonian formalism *à la* McCartor and Robertson. Because we do not have consistent pure space-like axial gauge quantization conditions, we have extrapolated the solution in the temporal gauge formulation and checked that it is also a solution in the axial gauge formulation. One conclusion to be drawn from the work involves the representation of the residual gauge fields: We have found that in the axial gauge we must choose the representation which specifies the ML prescription in order for the linear infrared divergences resulting from the operator $(\partial_-)^{-2}$ to be canceled. For this cancellation to work it is necessary for us to change integration variables to ones proportional to $n_+ = \sin 2\theta$. It follows from this that a pure space-like axial gauge formulation in the ordinary coordinates (this is the case of $\theta = \frac{\pi}{2}$ which gives us the Coulomb gauge) has to be defined as the limit $\theta \rightarrow \frac{\pi}{2} - 0$. Regulating the solution in this way involves information off the initial value surface ($x^0 = 0$), but in any event the solution includes auxiliary fields and the representation space is of indefinite metric.

In contrast, the case of $\theta = \frac{\pi}{4}$ (the light-cone axial gauge formulation) can be defined by simply setting $\theta = \frac{\pi}{4}$ (but the theory still has to be regulated by splitting the fermi products off the initial value surface).^{9,16)} This is because, by virtue of $n^2 = n_- = \cos 2\theta = 0$, we do not have the linear divergences resulting from $(\partial_-)^{-2}$ and from the square of the inverse of the hyperbolic Laplace operator except for the contact term in the most singular component of the gauge field propagator. We have also found that in order for the equal x^+ -time anticommutation relations to be satisfied in the axial gauges, we have to alter the representation of the $\tilde{\Sigma}$ from the temporal one to the axial one and make a modification to the overall normalization of the fermion field operators.

In contrast to the axial gauges, it seems that in the temporal gauges we can use either the representation of the residual gauge fields which specifies the ML prescription or the one which specifies the PV prescription. If we use the latter, then both components of the fermi field have vanishing vacuum expectation value.

Now that we have the operator solution in the axial gauge Schwinger model, we can

calculate commutation relations of the A_μ . It turns out that

$$[A_\mu(x), A_\nu(y)] = i\{-g_{\mu\nu}\Delta(x-y; m^2) + (n_\mu\partial_\nu + n_\nu\partial_\mu)\partial_- E(x-y) - n^2\partial_\mu\partial_\nu E(x-y)\}, \quad (4.1)$$

where $\Delta(x; m^2)$ is the commutator function of the free field of mass m and

$$E(x) = \frac{1}{\partial_-^2}\Delta(x; m^2) - \frac{x^-}{m^2 - n_- \partial_-^2}\delta(x^+) + \frac{2n_+\partial_+}{(m^2 - n_- \partial_-^2)^2}\delta(x^+). \quad (4.2)$$

It remains to be shown whether we can use (4.1) to obtain consistent pure space-like axial gauge quantization conditions which are needed to quantize interacting pure space-like axial gauge fields. We leave these tasks for subsequent studies.

Appendix A

— derivation of (3.16) —

In this appendix we give a detailed derivation of the x^+ -ordered gauge field propagator

$$D_{\mu\nu}(q) = \int d^2x \langle 0 | \{ \theta(x^+) A_\mu(x) A_\nu(0) + \theta(-x^+) A_\nu(0) A_\mu(x) \} | 0 \rangle e^{iq \cdot x}. \quad (A.1)$$

It is straightforward to show that the contributions from the $\tilde{\Sigma}$ and from the residual gauge fields are given, respectively, by

$$D_{++}^p(q) = \frac{i}{q^2 - m^2 + i\varepsilon} \frac{m^2}{q_-^2} = \frac{i}{q^2 - m^2 + i\varepsilon} \left(-g_{++} + \frac{2n_+q_+}{q_-} - n^2 \frac{q_+^2}{q_-^2} \right) - \frac{i}{q_-^2}, \quad (A.2)$$

$$\begin{aligned} D_{++}^g(q) = & \int_0^\infty dk_+ \left[\delta'(q_-) \frac{n_- k_+^2}{m^2 + n_- k_+^2} \left(\frac{i}{k_+ - q_+ - i\varepsilon} - \frac{i}{k_+ + q_+ - i\varepsilon} \right) \right. \\ & \left. + \delta(q_-) \frac{2n_+ k_+ m^2}{(n^2 k_+^2 + m^2)^2} \left(\frac{i}{k_+ - q_+ - i\varepsilon} + \frac{i}{k_+ + q_+ - i\varepsilon} \right) \right], \end{aligned} \quad (A.3)$$

Here, we have made use of the fact that $q^2 = n_- q_-^2 + 2n_+ q_+ q_- - n_- q_+^2$ and that $n^2 = n_-$. Note that the explicit x^- dependence gives rise to the factor $\delta'(q_-)$. Note also that there is no on-mass-shell condition for the residual gauge fields so that there remains a k_+ - integration. As a consequence, there arise singularities resulting from the inverse of the hyperbolic Laplace operator. It turns out that when we regularize the singularities by PV regularization, the integral on the first line of (A.3) is well-defined. In fact we can rewrite its integrand as a sum of simple poles:

$$\begin{aligned} \frac{n_- k_+^2}{m^2 + n_- k_+^2} \left(\frac{i}{k_+ - q_+ - i\varepsilon} - \frac{i}{k_+ + q_+ - i\varepsilon} \right) = & \frac{n_- q_+^2}{m^2 + n_- q_+^2} \left(\frac{i}{k_+ - q_+ - i\varepsilon} \right. \\ & \left. - \frac{i}{k_+ + q_+ - i\varepsilon} \right) - \frac{n_- a}{m^2 + n_- q_+^2} \left(\frac{i}{k_+ - a} - \frac{i}{k_+ + a} \right), \end{aligned} \quad (A.4)$$

where $a = \frac{m}{\sqrt{-n_-}}$. Direct calculation then gives

$$\int_0^\infty dk_+ \left(\frac{i}{k_+ - q_+ - i\varepsilon} - \frac{i}{k_+ + q_+ - i\varepsilon} \right) = -\pi \text{sgn}(q_+), \quad (\text{A}\cdot 5)$$

$$\text{P} \int_0^\infty dk_+ \left(\frac{1}{k_+ - a} - \frac{1}{k_+ + a} \right) = 0, \quad (\text{A}\cdot 6)$$

where $\text{sgn}(q_+)$ is obtained because the k_+ -integration region is limited to be $(0, \infty)$. On the other hand, the integral on the second line of (A·3) yields a linear divergence. We can see that by rewriting the integrand as a sum of simple and double poles:

$$\begin{aligned} & \frac{2n_+k_+m^2}{(m^2 + n_-k_+^2)^2} \left(\frac{i}{k_+ - q_+ - i\varepsilon} + \frac{i}{k_+ + q_+ - i\varepsilon} \right) = \frac{2n_+q_+m^2}{(m^2 + n_-q_+^2)^2} \\ & \times \left(\frac{i}{k_+ - q_+ - i\varepsilon} - \frac{i}{k_+ + q_+ - i\varepsilon} \right) + n_+a \frac{n_-q_+^2 - m^2}{(m^2 + n_-q_+^2)^2} \left(\frac{i}{k_+ - a} - \frac{i}{k_+ + a} \right) \\ & - \frac{n_+}{n_-} \frac{m^2}{m^2 + n_-q_+^2} \left(\frac{i}{(k_+ - a)^2} + \frac{i}{(k_+ + a)^2} \right). \end{aligned} \quad (\text{A}\cdot 7)$$

We can evaluate the integrations of the first and second terms on the right hand with the help of (A·5) and (A·6). However, we cannot regularize a linear divergence resulting from the double pole by the PV prescription. We show below that this linear divergence cancels one resulting from the factor $(q_-^2)^{-1}$ of D_{++}^p in (A·2). For later convenience we rewrite the linearly diverging integration in the form

$$\text{P} \int_0^\infty dk_+ \left(\frac{n_+}{(k_+ - a)^2} + \frac{n_+}{(k_+ + a)^2} \right) = \text{P} \int_{-\infty}^\infty dk_+ \frac{n_+}{(k_+ - a)^2} = - \int_{-\infty}^\infty dq_- \frac{n_-}{q_-^2}, \quad (\text{A}\cdot 8)$$

where we have changed the integration variable from k_+ to $q_- = \frac{-n_-}{n_+}(k_+ - a)$. Substituting (A·5), (A·6) and (A·8) into (A·3) yields

$$\begin{aligned} D_{++}^q(q) = & \frac{i}{q^2 - m^2 + i\varepsilon} \left(-in^2q_+^2\pi\text{sgn}(q_+)\delta'(q_-) - 2in_+q_+\delta(q_-)\pi\text{sgn}(q_+) \right. \\ & \left. - m^2\delta(q_-)\int_{-\infty}^\infty dq_- \frac{1}{q_-^2} \right), \end{aligned} \quad (\text{A}\cdot 9)$$

where we have made use of the identity

$$\frac{1}{q^2 - m^2 + i\varepsilon} \delta'(q_-) = -\frac{1}{m^2 + n_-q_+^2} \delta'(q_-) + \frac{2n_+q_+}{(m^2 + n_-q_+^2)^2} \delta(q_-). \quad (\text{A}\cdot 10)$$

Thus, for the sum of (A·2) and (A·9) we obtain

$$\begin{aligned} D_{++}(q) = & \frac{i}{q^2 - m^2 + i\varepsilon} \left(-g_{++} + \frac{2n_+q_+}{q_- + i\varepsilon\text{sgn}(q_+)} - in^2q_+^2\pi\text{sgn}(q_+)\delta'(q_-) \right. \\ & \left. - n^2q_+^2\left(\frac{1}{q_-^2} - \delta(q_-)\int_{-\infty}^\infty dq_- \frac{1}{q_-^2}\right) - i\left(\frac{1}{q_-^2} - \delta(q_-)\int_{-\infty}^\infty dq_- \frac{1}{q_-^2}\right), \right. \end{aligned} \quad (\text{A}\cdot 11)$$

where we have made use of the identity:

$$\delta(q_-) \frac{-m^2}{q_-^2 - m^2 + i\varepsilon} = \delta(q_-) \left(1 + \frac{n^2 q_+^2}{q_-^2 - m^2 + i\varepsilon} \right). \quad (\text{A}\cdot 12)$$

Now we show that the term $\frac{1}{q_-^2} - \delta(q_-) \int_{-\infty}^{\infty} dq_- \frac{1}{q_-^2}$ does not give rise to any divergences when we calculate $D_{++}(x)$ by substituting (A.11) into (A.1). To show this we consider the integration:

$$\begin{aligned} & \int_{-\infty}^{\infty} dq_- \frac{1}{q_-^2 - m^2 + i\varepsilon} \left(\frac{1}{q_-^2} - \delta(q_-) \int_{-\infty}^{\infty} dq_- \frac{1}{q_-^2} \right) e^{-iq_- x^-} \\ &= \int_{-\infty}^{\infty} dq_- \frac{e^{-iq_- x^-}}{q_-^2 (q_-^2 - m^2 + i\varepsilon)} + \frac{1}{m^2 + n_- q_+^2} \int_{-\infty}^{\infty} dq_- \frac{1}{q_-^2}. \end{aligned} \quad (\text{A}\cdot 13)$$

We can rewrite the second line further as

$$\begin{aligned} & \int_{-\infty}^{\infty} dq_- \frac{e^{-iq_- x^-} - 1}{q_-^2} \frac{1}{q_-^2 - m^2 + i\varepsilon} + \int_{-\infty}^{\infty} dq_- \frac{1}{q_-^2} \left(\frac{1}{m^2 + n_- q_+^2} + \frac{1}{q_-^2 - m^2 + i\varepsilon} \right) \\ &= \int_{-\infty}^{\infty} dq_- \left(\frac{e^{-iq_- x^-} - 1}{q_-^2} \frac{1}{q_-^2 - m^2 + i\varepsilon} + \frac{2n_+ q_+ + n_- q_-}{q_- (m^2 + n_- q_+^2) (q_-^2 - m^2 + i\varepsilon)} \right). \end{aligned} \quad (\text{A}\cdot 14)$$

We see that the last integrals diverge at most logarithmically; but logarithmic divergences can be regularized by the PV prescription so there arise no divergences in (A.14). This verifies that the following identity holds:

$$\begin{aligned} & \frac{1}{q_-^2} + i\pi \text{sgn}(q_+) \delta'(q_-) - \delta(q_-) \int_{-\infty}^{\infty} dq_- \frac{1}{q_-^2} \\ &= \text{Pf} \frac{1}{q_-^2} + i\pi \text{sgn}(q_+) \delta'(q_-) = \frac{1}{(q_- + i\varepsilon \text{sgn}(q_+))^2} \end{aligned} \quad (\text{A}\cdot 15)$$

where Pf denotes Hadamard's finite part. It should be also noted that

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} dq_- \left(\delta(q_-) \int_{-\infty}^{\infty} dq_- \frac{1}{q_-^2} - \frac{1}{q_-^2} \right) e^{-q_- x^-} = \frac{1}{\pi} \int_0^{\infty} dq_- \frac{1 - \cos q_- x^-}{q_-^2} = \frac{|x^-|}{2}, \quad (\text{A}\cdot 16)$$

where the Fourier transform of the last term is $-\frac{1}{2} \left(\frac{1}{(q_- + i\varepsilon)^2} + \frac{1}{(q_- - i\varepsilon)^2} \right)$. Thus we obtain

$$\begin{aligned} D_{++}(q) &= \frac{i}{q^2 - m^2 + i\varepsilon} \left(-g_{++} + \frac{2n_+ q_+}{q_- + i\varepsilon \text{sgn}(q_+)} - \frac{n^2 q_+^2}{(q_- + i\varepsilon \text{sgn}(q_+))^2} \right) \\ &\quad - \delta_{\mu+} \delta_{\nu+} \frac{i}{2} \left(\frac{1}{(q_- + i\varepsilon)^2} + \frac{1}{(q_- - i\varepsilon)^2} \right). \end{aligned} \quad (\text{A}\cdot 17)$$

This completes showing that, due to the residual gauge fields, the linear divergence resulting from the ∂_+^{-1} is eliminated from the x^+ - ordered gauge field propagator.

References

- 1) Yu. L. Dokshitzer, D. J. D'Yakonov, and S. Y. Troyan, Phys. Rep. **58** (1980), 270.
- 2) L. Susskind, Phys. Rev. **165** (1968), 1535.
K. Bardakci and M. B. Halpern, Phys. Rev. **176** (1968), 1686.
- 3) S. J. Brodsky, H. C. Pauli and S. S. Pinsky, Phys. Rep. **301** (1998), 299.
- 4) N. Nakanishi, Phys. Lett. **131B** (1983), 381.
N. Nakanishi, *Quantum Electrodynamics*, ed. T. Kinoshita (World Scientific, Singapore, 1990), p. 36.
- 5) D. M. Capper, J. J. Dulwich, and M. J. Litvak, Nucl. Phys. **241B** (1984), 463.
S. Mandelstam, Nucl. Phys. **B213** (1983), 149.
G. Leibbrandt, Phys. Rev. **D29** (1984), 1699.
- 6) A. Bassetto, M. Dalbosco, I. Lazziera and R. Soldati, Phys. Rev. **D31** (1985), 2012.
A. Bassetto, G. Nardelli and R. Soldati, *Yang-Mills Theories in Algebraic Non-Covariant Gauges* (World Scientific, Singapore, 1991).
- 7) M. Morara and R. Soldati, Phys. Rev. **D58** (1998), 105011.
- 8) G. McCartor and D.G. Robertson, Z. Phys. **C62** (1994), 349; **C68** (1995), 345.
- 9) Y. Nakawaki and G. McCartor, Prog. Theor. Phys. **103** (2000), 161.
- 10) I. Lazzizzera, Phys. Lett. **210B** (1988), 188; Nuovo Cim. **102A** (1989), 1385.
- 11) P. V. Landshoff and P. van Nieuwenhuizen, Phys. Rev. **D50** (1994), 4157.
- 12) P. A. M. Dirac, *Lectures on Quantum Mechanics*, Belfer Graduate School of Science-Yeshiva University (Academic Press, New York, 1964).
A. Hanson, T. Regge and C. Teitelboim, *Constrained Hamiltonian Systems* (Accad. Naz. dei Lincei, Rome, 1976).
- 13) Y. Nakawaki and G. McCartor, Prog. Theor. Phys. **102** (1999), 149.
- 14) K. Hornbostel, Phys. Rev. **D45** (1992), 3781; E. Prokhvatilov and V. Franke, Yad. Fiz. **49** (1989), 1109; F. Lenz, M. Thies, S. Levit, and D. Yazaki, Ann. Phys. **208** (1991), 1.
- 15) M. Abe and S. Miyake, Phys. Lett. **76B** (1978), 76; Prog. Theor. Phys. **59** (1978), 2037.
- 16) G. McCartor, Phys. Rev. **D60** (1999), 105004; **hep-th/0008040**.
- 17) B. Lautrup, Mat. Fys. Medd. Dan. Vid. Selsk. **35**(1967), No.11.
N. Nakanishi, Prog. Theor. Phys. Suppl. No.51 (1972), 1.
- 18) Y. Nakawaki, Prog. Theor. Phys. **72** (1984), 152.
- 19) K. Haller, Phys. Rev. **D42** (1990), 2095.
- 20) G. McCartor, Z. Phys. **C41** (1988), 271.